

Comparison of Long-Acting Lenacapavir Phase 2/3 Regimen vs Simplified Regimen Using Population-PK Analysis and Simulation



Naveed A. Shaik,¹ Francesco Bellanti,² Kishore Pollireddy,² Vamshi Jogiraju,¹ Craig Comisar,² Sandhya Girish,¹ Martin Rhee,¹ Ramesh Palaparthi,¹ Renu Singh¹

¹Gilead Sciences, Inc., Foster City, California, USA; ²Certara, Inc., Princeton, New Jersey, USA

Key Findings

- Lenacapavir (LEN) is approved for the treatment of multidrug-resistant HIV-1 in combination with other antiretrovirals for heavily treatment-experienced people with HIV (HTE PWH)
- Two LEN dosing regimens are approved in the USA:
 - Phase 2/3 regimen: Days 1 and 2: 600 mg orally; Day 8: 300 mg orally then subcutaneous (SC) LEN dosing (927 mg every 6 months [Q6M]) starting on Day 15
 - Simplified regimen: Day 1: 600 mg orally and 927 mg SC injection; Day 2: 600 mg orally; then SC LEN dosing with 927 mg SC Q6M
- Pharmacokinetic (PK) data in HTE PWH are currently not available for the simplified regimen
- Simulation using a LEN PopPK model indicated that the simplified regimen was comparable with the Phase 2/3 regimen in HTE PWH: at steady state the accumulation ratio was 1.2-fold for both regimens
- Thus, no differences in the efficacy and safety of LEN in HTE PWH are expected between the two regimens


Conclusions

- Simulated LEN PK for the first SC dose and at steady state for the Phase 2/3 and simplified regimens were comparable across all exposure metrics with steady-state LEN concentrations being identical in HTE PWH
- No differences in the safety and efficacy of LEN are expected based on the PK similarity of the Phase 2/3 and simplified regimens
- These data support the use of the simplified regimen for ongoing LEN treatment and prevention studies

Background

- LEN, a potent first-in-class capsid inhibitor, is approved for the treatment of multidrug-resistant HIV-1 in combination with other antiretrovirals for HTE PWH^{1,2}
- LEN has shown near maximum antiviral activity when the lower bound of the 90% confidence interval (CI) of mean trough concentration (C_{trough}) is maintained above 15.5 ng/mL,³ which is the inhibitory quotient-4 (IQ4; ≥ 4 -fold greater than the *in vitro* protein-adjusted 95% effective concentration in MT-4 cells)⁴
- In the ongoing pivotal Phase 2/3 study (CAPELLA, NCT04150068⁵), participants received oral LEN loading doses (Days 1 and 2: 600 mg; Day 8: 300 mg) then SC LEN dosing (927 mg Q6M) starting on Day 15 (Phase 2/3 regimen; **Figure 1**)
- This Phase 2/3 regimen and a simplified regimen (Day 1: 600mg orally and 927 mg SC injection; Day 2: 600 mg orally; then SC LEN dosing with 927 mg SC Q6M; **Figure 1**) were recently approved by the FDA¹
- In a healthy volunteer study, LEN plasma concentrations were comparable between the Phase 2/3 regimen and the simplified regimen for 6 months after the first SC dose⁶
- PK data in HTE PWH are currently not available for the simplified regimen

Figure 1. Phase 2/3 and simplified LEN dosing regimens



d, day; LEN, lenacapavir; SC, subcutaneous, wk, week.

Objective

- To compare the simulated steady-state LEN exposure metrics between the Phase 2/3 and simplified regimens in HTE PWH

Methods

- LEN plasma concentrations were simulated using a previously developed 2-compartment population-PK (PopPK) model with 1st-order process for oral absorption, and parallel 1st-order and transit compartments for SC absorption and linear elimination⁷
- Using the PopPK model, plasma concentrations were simulated with both the Phase 2/3 and the simplified regimens (**Figure 2**), and LEN exposure metrics were compared (**Figure 3**)
- Key exposure metrics were the area under the concentration-time curve over the dosing interval (AUC_{tau}), maximum concentration of LEN (C_{max}), and C_{trough}
- External validation was conducted using observed data for the simplified regimen in healthy participants (Phase I study, GS-US-200-5709, Cohort 2)

Results

PK Simulations

- Simulated LEN exposure in HTE PWH for the Phase 2/3 and simplified regimens are shown in **Table 1** for the first dose and at steady-state
- The simplified regimen was comparable with the Phase 2/3 regimen up to 6 months after the first SC dose (**Table 1**)
- Exposure metrics (AUC_{tau} , C_{max} , C_{trough}) were identical at steady-state for both regimens (**Table 1**)
- Steady-state was achieved by the second SC dose with 1.2-fold accumulation for both regimens (**Figure 2**)
- As shown in **Figure 3C**, the lower bounds of the 90% CIs of simulated mean LEN C_{trough} were found to be consistently above the IQ4 threshold of 15.5 ng/mL for both regimens over the treatment duration

External validation


- External validation indicated that the PopPK model captured the simplified regimen data adequately (**Figure 4**)
- As C_{max} is driven by SC administration, both regimens resulted in similar C_{max} over the duration of treatment (**Figure 4**), thus there are no safety concerns

Table 1. Simulated LEN exposure metrics for Phase 2/3 and simplified regimens of LEN in HTE PWH

Parameter, Mean (%CV)	Phase 2/3 regimen			Simplified regimen		
	Days 1–15	Day 15 – End of month 6	Steady state	Days 1–15	Day 15 – End of month 6	Steady state
AUC_{tau} , h•ng/mL	15,600 (52.9)	250,000 (66.6)	300,000 (68.5)	18,800 (53.6)	238,000 (67.5)	300,000 (68.5)
C_{max} , ng/mL	69.6 (56.0)	87 (71.8)	97.2 (70.3)	80.1 (55.7)	87.1 (71.9)	97.2 (70.3)
C_{trough} , ng/mL	35.9 (56.8)	32.7 (88)	36.2 (90.6)	49 (57.9)	32.7 (88)	36.2 (90.6)


AUC_{tau} , area under curve over dosing interval; C_{max} , maximum concentration; C_{trough} , trough concentration; HTE, heavily treatment-experienced; LEN, lenacapavir; PWH, people living with HIV; %CV, percentage coefficient of variation, simulations conducted using approximately 40,000 virtual patients

Figure 2. Simulated plasma LEN concentration-time curves with multiple dosing for Phase 2/3 and simplified regimens in HTE PWH




CI, confidence intervals; HTE, heavily treatment-experienced; LEN, lenacapavir; IQ4, inhibitory quotient-4; PWH, people with HIV.

Figure 3. Simulated A) LEN AUC_{tau} , B) LEN C_{max} , and C) LEN C_{trough} for the Phase 2/3 and simplified regimens in HTE PWH over the treatment duration for the first dose



AUC_{tau} , area under curve over dosing interval; C_{max} , maximum concentration; C_{trough} , trough concentration; HTE, heavily treatment-experienced; K, thousand; LEN, lenacapavir; PWH, people with HIV.

Figure 4. Prediction-corrected visual predictive check plots – External validation of simplified LEN regimen



Prediction-corrected visual predictive check plots show medians (solid red lines) and spreads (5th-95th percentiles [dashed red lines]) of observed data in all participants; red areas are 95% CIs of simulated medians and blue areas are 95% CIs of simulated 5th and 95th percentiles; black circles are individual observed data corrected by model predictions; orange dashes indicate boundaries of visual predictive check bins. CI, confidence interval; d, day; LEN, lenacapavir; PK, pharmacokinetic; wk, week.

References: 1. Sunlenca US PI SmPC 2022, available on https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/215973s000lbl.pdf last accessed 10 July 2023; 2. Sunlenca EU SmPC 2022, available on https://www.ema.europa.eu/en/documents/product-information/sunlenca-epar-product-information_en.pdf last accessed 10 July 2023; 3. Shaik N et al. Poster PESUB23 presented at AIDS 2022; 4. Link JO, et al. *Nature* 2020;584:614–618/ 5. Segal-Maurer S, et al. *N Engl J Med* 2022;386:1793–1803; 6. Jogiraju V et al. Poster PESUB22 presented at AIDS 2022; 7. Shaik N et al. Poster EPB174 presented at AIDS 2022.

Acknowledgments: These analyses were conducted by Certara, Inc., and were funded by Gilead Sciences, Inc. Medical writing support was provided by Jackie Phillipson of Ashfield MedComms, an Inizio company, Macclesfield, UK, and was funded by Gilead Sciences, Inc.

Disclosures: NA Shaik, V Jogiraju, S Girish, M Rhee, R Palaparthi, R Singh: employees and shareholders of Gilead Sciences, Inc. F Bellanti, K Pollireddy, C Comisar: employees and shareholders of Certara, Inc.